



Is JELMYTO right for you? Your doctor discussion guide


A diagnosis of low-grade upper tract urothelial carcinoma (UTUC) may seem daunting, but by asking your doctor the right questions you can take an active role in your treatment path.

Because you have **low-grade UTUC**, JELMYTO may be an option for you. This guide will help you learn more about your condition and treatment with JELMYTO.

Use the drawing below to talk to your doctor about your UTUC.

Where is my tumor/where are my tumors located?

Have your doctor draw on the diagram



Other information about my UTUC:

Approved Use for JELMYTO

JELMYTO® is a prescription medicine used to treat adults with a type of cancer of the lining of the upper urinary tract including the kidney called low-grade Upper Tract Urothelial Cancer (LG-UTUC).

Please [click here](#) for JELMYTO Full Prescribing Information, including the Patient Information, for additional information.

Here are some other questions you can ask your doctor to find out if JELMYTO is the right treatment for you.

Am I a candidate for JELMYTO?

What makes JELMYTO different from other treatment options?

Has JELMYTO helped people with my type of cancer?

What can I expect when I start taking JELMYTO?

How often will I get JELMYTO?

What kind of recovery comes with JELMYTO?

What are the possible side effects of JELMYTO treatment?

Please write down any other notes or questions here:

Important Safety Information

You should not receive JELMYTO if you have a hole or tear (perforation) of your bladder or upper urinary tract.

Before receiving JELMYTO, tell your healthcare provider about all your medical conditions, including if you:

- are pregnant or plan to become pregnant. JELMYTO can harm your unborn baby. You should not become pregnant during treatment with JELMYTO. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with JELMYTO.

Females who are able to become pregnant: You should use effective birth control (contraception) during treatment with JELMYTO and for 6 months after the last dose.

Please [click here](#) for JELMYTO Full Prescribing Information, including the Patient Information, for additional information.



“I’m ready to ask the right questions so I can fight my UTUC.”

Important Safety Information (cont’d)

Males being treated with JELMYTO: If you have a female partner who is able to become pregnant, you should use effective birth control (contraception) during treatment with JELMYTO and for 3 months after the last dose.

- are breastfeeding or plan to breastfeed. It is not known if JELMYTO passes into your breast milk. Do not breastfeed during treatment with JELMYTO and for 1 week after the last dose.
- **Tell your healthcare provider if you take water pills (diuretic).**

How will I receive JELMYTO?

- Your healthcare provider will tell you to take a medicine called sodium bicarbonate before each JELMYTO treatment.
- You will receive your JELMYTO dose from your healthcare provider 1 time a week for 6 weeks. It is important that you receive all 6 doses of JELMYTO according to your healthcare provider’s instructions. If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment. Your healthcare provider may recommend up to an additional 11 monthly doses.
- JELMYTO is given to your kidney through a tube called a catheter.
- During treatment with JELMYTO, your healthcare provider may tell you to take additional medicines or change how you take your current medicines.

After receiving JELMYTO:

- JELMYTO may cause your urine color to change to a violet to blue color. Avoid contact between your skin and urine for at least 6 hours.
- To urinate, **males and females should sit** on a toilet and flush the toilet several times after you use it. After going to the bathroom, wash your hands, your inner thighs, and genital area well with soap and water.
- Clothing that comes in contact with urine should be washed right away and washed separately from other clothing.

JELMYTO may cause serious side effects, including:

- **Swelling and narrowing of the tube that carries urine from the kidney to the bladder (ureteric obstruction).** If you develop swelling and narrowing, and to protect your kidney from damage, your healthcare provider may recommend the placement of a small plastic tube (stent) in the ureter to help the kidney drain. Tell your healthcare provider right away if you develop side pain or fever during treatment with JELMYTO.
- **Bone marrow problems.** JELMYTO can affect your bone marrow and can cause a decrease in your white blood cell, red blood cell, and platelet counts. Your healthcare provider will do blood tests prior to each treatment to check your blood cell counts during treatment with JELMYTO. Your healthcare provider may need to temporarily or permanently stop JELMYTO if you develop bone marrow problems during treatment with JELMYTO.

The most common side effects of JELMYTO include: urinary tract infection, blood in your urine, side pain, nausea, trouble with urination, kidney problems, vomiting, tiredness, stomach (abdomen) pain.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to UroGen Pharma at 1-855-987-6436.

Ask your doctor about JELMYTO, a nonsurgical option for low-grade UTUC

Please [click here](#) for JELMYTO Full Prescribing Information, including the Patient Information, for additional information.



JELMYTO® and UroGen® are registered trademarks of UroGen Pharma, Ltd.
© 2021 UroGen Pharma, Inc. All rights reserved. US-JEL-00306 02/21

